Differentiation (2000) 65:301–306
© Springer-Verlag 2000

SUBJECT INDEX Volume 65 (1999/2000)

Adipocyte differentiation
Inoue-Murayama M., Sugimoto Y., Niimi Y., Aso H.
Type XVIII collagen is newly transcribed during bovine adipogenesis, 65:281–285

Adipogenesis
Inoue-Murayama M., Sugimoto Y., Niimi Y., Aso H.
Type XVIII collagen is newly transcribed during bovine adipogenesis, 65:281–285

Alopecia
Wojcik S.M., Imakado S., Seki T., Longley M.A., Petherbridge L., Bundman D.S., Bickenbach J.R., Rothnagel J.A., Roop D.R.
Expression of *MK6a* dominant-negative and C-terminal mutant transgenes in mice has distinct phenotypic consequences in the epidermis and hair follicle, 65:97–112

Axis defects
Gong S.-G., Kiba A.
The role of Xmsx-2 in the anterior-posterior patterning of the mesoderm in *Xenopus laevis*, 65:131–140

Axonformation
Leake D., Asch W.S., Canger A.K., Schechter N.
Gefiltin in zebrafish embryos: sequential gene expression of two neurofilament proteins in retinal ganglion cells, 65:181–189

Basonuclin
Tseng H., Matsuzaki K., Lavker R.M.
Basonuclin in murine corneal and lens epithelia correlates with cellular maturation and proliferative ability, 65:221–227

Bladder epithelium
Liang F., Kachar B., Ding M., Zhai Z., Wu X.-R., Sun T.-T.
Urothelial hinge as a higly specialized membrane: detergent-insolubility, urohingin association, and in vitro formation, 65:59–69

Aboseif S., El-Sakka A., Young P., Cunha G.
Mesenchymal reprogramming of adult human epithelial differentiation, 65:113–118

Bone marrow-derived mast cells
Fehlner-Gardiner C.C., Cao H., Jackson-Boeters L., Nakamura T., Elliott B.E., Uniyal S., Chan B.M.C.
Characterization of a functional relationship between hepatocyte growth factor and mouse bone marrow-derived mast cells, 65:27–42

Calcium
Joffroy S., Letellier T., Rossignol R., Malgat M., Delage J.-P., Powell J.A., Mazat J.-P., Koenig J.
Modification of mitochondrial metabolism in fibroblasts from mice with a skeletal muscle mutation (muscular dysgenesis): Evidence of embryonic communication between myoblasts and fibroblasts, 65:261–270

CD14
Pujol B.F., Lucibello F.C., Gehling U.M., Lindemann K., Weidner N., Zuzarte M.-L., Adamkiewicz J., Elsässer H.-P., Müller R., Havemann K.
Endothelioid cells derived from human CD14 positive monocytes, 65:287–300

Cell migration
Fehlner-Gardiner C.C., Cao H., Jackson-Boeters L., Nakamura T., Elliott B.E., Uniyal S., Chan B.M.C.
Characterization of a functional relationship between hepatocyte growth factor and mouse bone marrow-derived mast cells, 65:27–42

Chicken stomach
Koike T., Yasugi S.
In vitro analysis of mesenchymal influences on the differentiation of stomach epithelial cells of the chicken embryo. 65:13–25

Collagen
Inoue-Murayama M., Sugimoto Y., Niimi Y., Aso H.
Type XVIII collagen is newly transcribed during bovine adipogenesis, 65:281–285

Cornea
Mizuno N., Mochii M., Yamamoto T.S., Takahashi T.C., Eguchi G., Okada T.S.
Pax-6 and *Prox 1* expression during lens regeneration from *Cynops* iris and *Xenopus* cornea: evidence for a genetic program common to embryonic lens development, 65:141–149

Cynops
Mizuno N., Mochii M., Yamamoto TS., Takahashi T.C., Eguchi G., Okada T.S.
Pax-6 and *Prox 1* expression during lens regeneration from *Cynops* iris and *Xenopus* cornea: evidence for a genetic program common to embryonic lens development, 65:141–149

Cytokeratins
Benazzouz A., Duprey P.
The vimentin promoter as a tool to analyze the early events of retinoic acid-induced differentiation of cultured embryonal carcinoma cells, 65:171–180

Rothnagel J.A., Seki T., Ogo M., Longley M.A., Wojcik S.M., Bundman D.S., Bickenbach J.R., Roop D.R.
The mouse keratin 6 isoforms are differentially expressed in the hair follicle, footpad, tongue and activated epidermis, 65:119–130

Desmin
Loh S.-H., Chan W.-T., Gong Z., Lim T.-M., Chua K.-L.
Characterization of a zebrafish (*Danio rerio*) desmin cDNA: an early molecular marker of myogenesis, 65:247–254

Dictyostelium
Agarwal A.K., Parrish S.N., Blumberg D.D.
Ribosomal protein gene expression is cell type specific during development in *Dictyostelium discoideum*, 65:73–88

eIF4E
Fahrenkrug S.C., Dahlquist M.O., Clark K.J., Hackett P.B.
Dynamic and tissue-specific expression of eIF4E during zebrafish embryogenesis, 65:191–201

Embryonal carcinoma cells
Benazzouz A., Duprey P.
The vimentin promoter as a tool to analyze the early events of retinoic acid-induced differentiation of cultured embryonal carcinoma cells, 65:171–180

Tomotsune D., Takihara Y., Berger J., Duhl D., Joo S., Kyba M., Shirai M., Ohta H., Matsuda Y., Honda B.M., Simon J., Shimada K., Brock H.W., Randazzo F.
A novel member of murine Polycomb-group proteins, Sex comb on midleg homolog protein, is highly conserved, and interacts with RAE28/mph1 in vitro, 65:229–239

Pratt M.A.C., Crippen C.A., Ménard M.
Spontaneous retinoic acid receptor β2 expression during mesoderm differentiation of P19 murine embryonal carcinoma cells, 65:271–279

Embryonic chicken pepsinogen (ECPg)
Koike T., Yasugi S.
In vitro analysis of mesenchymal influences on the differentiation of stomach epithelial cells of the chicken embryo. 65:13–25

Endothel
Pujol B.F., Lucibello F.C., Gehling U.M., Lindemann K., Weidner N., Zuzarte M.-L., Adamkiewicz J., Elsässer H.-P., Müller R., Havemann K.
Endothelioid cells derived from human CD14 positive monocytes, 65:287–300

Epidermis
Wojcik S.M., Imakado S., Seki T., Longley M.A., Petherbridge L., Bundman D.S., Bickenbach J.R., Rothnagel J.A., Roop D.R.
Expression of *MK6a* dominant-negative and C-terminal mutant transgenes in mice has distinct phenotypic consequences in the epidermis and hair follicle, 65:97–112

Rothnagel J.A., Seki T., Ogo M., Longley M.A., Wojcik S.M., Bundman D.S., Bickenbach J.R., Roop D.R.
The mouse keratin 6 isoforms are differentially expressed in the hair follicle, footpad, tongue and activated epidermis, 65:119–130

Epithelia
Gulyás M., Dobra K., Hjerpe A.
Expression of genes coding for proteoglycans and Wilms' tumour susceptibility gene 1 (*WT1*) by variously differentiated benign human mesothelial cells, 65:89–96

Tseng H., Matsuzaki K., Lavker R.M.
Basonuclin in murine corneal and lens epithelia correlates with cellular maturation and proliferative ability, 65:221–227

Epithelial-mesenchymal interaction
Koike T., Yasugi S.
In Vitro analysis of mesenchymal influences on the differentiation of stomach epithelial cells of the chicken embryo. 65:13–25

Lichter-Konecki U., Moter S.E., Krawisz B.R., Schlotter M., Hipke C., Konecki D.S.
Expression patterns of murine lysosome-associated membrane protein 2 (Lamp-2) transcripts during morphogenesis, 65:43–58

Aboseif S., El-Sakka A., Young P., Cunha G.
Mesenchymal reprogramming of adult human epithelial differentiation, 65:113–118

Erythropoiesis
Fahrenkrug S.C., Dahlquist M.O., Clark K.J., Hackett P.B.
Dynamic and tissue-specific expression of eIF4E during zebrafish embryogenesis, 65:191–201

Extracellular matrix
Lichter-Konecki U., Moter S.E., Krawisz B.R., Schlotter M., Hipke C., Konecki D.S.
Expression patterns of murine lysosome-associated membrane protein 2 (Lamp-2) transcripts during morphogenesis, 65:43–58

Fibroblast growth factor
Koike T., Yasugi S.
In vitro analysis of mesenchymal influences on the differentiation of stomach epithelial cells of the chicken embryo. 65:13–25

Düsterhöft S., Pette D.
Evidence that acidic fibroblast growth factor promotes maturation of rat satellite-cell-derived myotubes in vitro, 65:161–169

Düsterhöft S., Putman C.T., Pette D.
Changes in FGF and FGF-receptor expression in low-frequency-stimulated rat muscles and rat satellite cell cultures, 65:203–208

Fibroblast-like cells
Gulyás M., Dobra K., Hjerpe A.
Expression of genes coding for proteoglycans and Wilms' tumour susceptibility gene 1 (*WT1*) by variously differentiated benign human mesothelial cells, 65:89–96

Joffroy S., Letellier T., Rossignol R., Malgat M., Delage J.-P., Powell J.A., Mazat J.-P., Koenig J.
Modification of mitochondrial metabolism in fibroblasts from mice with a skeletal muscle mutation (muscular dysgenesis): Evidence of embryonic communication between myoblasts and fibroblasts, 65:261–270

Gap junctions
Liptau H., Viebahn C.
Expression patterns of gap junctional proteins connexin 32 and 43 suggest new communication compartments in the gastrulating rabbit embryo, 65:209–219

Gastrulation
Fahrenkrug S.C., Dahlquist M.O., Clark K.J., Hackett P.B.
Dynamic and tissue-specific expression of eIF4E during zebrafish embryogenesis, 65:191–201

Liptau H., Viebahn C.
Expression patterns of gap junctional proteins connexin 32 and 43 suggest new communication compartments in the gastrulating rabbit embryo, 65:209–219

Germ layers
Liptau H., Viebahn C.
Expression patterns of gap junctional proteins connexin 32 and 43 suggest new communication compartments in the gastrulating rabbit embryo, 65:209–219

Gefiltin
Leake D., Asch W.S., Canger A.K., Schechter N.
Gefiltin in zebrafish embryos: sequential gene expression of two neurofilament proteins in retinal ganglion cells, 65:181–189

Hair follicle
Wojcik S.M., Imakado S., Seki T., Longley M.A., Petherbridge L., Bundman D.S., Bickenbach J.R., Rothnagel J.A., Roop D.R.
Expression of *MK6a* dominant-negative and C-terminal mutant transgenes in mice has distinct phenotypic consequences in the epidermis and hair follicle, 65:97–112

Rothnagel J.A., Seki T., Ogo M., Longley M.A., Wojcik S.M., Bundman D.S., Bickenbach J.R., Roop D.R.
The mouse keratin 6 isoforms are differentially expressed in the hair follicle, footpad, tongue and activated epidermis, 65:119–130

Hepatocyte growth factor
Fehlner-Gardiner C.C., Cao H., Jackson-Boeters L., Nakamura T., Elliott B.E., Uniyal S., Chan B.M.C.
Characterization of a functional relationship between hepatocyte growth factor and mouse bone marrow-derived mast cells, 65:27–42

Hindbrain
Essner J.J., Johnson R.G., Hackett P.B Jr.
Overexpression of thyroid hormone receptor α1 during zebrafish embryogenesis disrupts hindbrain patterning and implicates retinoic acid receptors in the control of *hox* gene expression, 65:1–11

Histone H1
Flickinger R.A.
Induction of tumor cell differentiation, 65:241–245

Histones
Flickinger R.A.
Induction of tumor cell differentiation, 65:241–245

Hox genes
Essner J.J., Johnson R.G., Hackett P.B Jr.
Overexpression of thyroid hormone receptor α1 during zebrafish embryogenesis disrupts hindbrain patterning and implicates retinoic acid receptors in the control of *hox* gene expression, 65:1–11

Mizuno N., Mochii M., Yamamoto T.S., Takahashi T.C., Eguchi G., Okada T.S.
Pax-6 and *Prox 1* expression during lens regeneration from *Cynops* iris and *Xenopus* cornea: evidence for a genetic program common to embryonic lens development, 65:141–149

Tomotsune D., Takihara Y., Berger J., Duhl D., Joo S., Kyba M., Shirai M., Ohta H., Matsuda Y., Honda B.M., Simon J., Shimada K., Brock H.W., Randazzo F.
A novel member of murine Polycomb-group proteins, Sex comb on midleg homolog protein, is highly conserved, and interacts with RAE28/mph1 in vitro, 65:229–239

Immunohistochemistry
Liptau H., Viebahn C.
Expression patterns of gap junctional proteins connexin 32 and 43 suggest new communication compartments in the gastrulating rabbit embryo, 65:209–219

In-situ hybridization
Loh S.-H., Chan W.-T., Gong Z., Lim T.-M., Chua K.-L.
Characterization of a zebrafish (*Danio rerio*) desmin cDNA: an early molecular marker of myogenesis, 65:247–254

β1 integrins
Fehlner-Gardiner C.C., Cao H., Jackson-Boeters L., Nakamura T., Elliott B.E., Uniyal S., Chan B.M.C.
Characterization of a functional relationship between hepatocyte growth factor and mouse bone marrow-derived mast cells, 65:27–42

Intercellular communication
Liptau H., Viebahn C.
Expression patterns of gap junctional proteins connexin 32 and 43 suggest new communication compartments in the gastrulating rabbit embryo, 65:209–219

Intermediate filaments
Wojcik S.M., Imakado S., Seki T., Longley M.A., Petherbridge L., Bundman D.S., Bickenbach J.R., Rothnagel J.A., Roop D.R.
Expression of *MK6a* dominant-negative and C-terminal mutant transgenes in mice has distinct phenotypic consequences in the epidermis and hair follicle, 65:97–112

Rothnagel J.A., Seki T., Ogo M., Longley M.A., Wojcik S.M., Bundman D.S., Bickenbach J.R., Roop D.R.
The mouse keratin 6 isoforms are differentially expressed in the hair follicle, footpad, tongue and activated epidermis, 65:119–130

Benazzouz A., Duprey P.
The vimentin promoter as a tool to analyze the early events of retinoic acid-induced differentiation of cultured embryonal carcinoma cells, 65:171–180

Leake D., Asch W.S., Canger A.K., Schechter N.
Gefiltin in zebrafish embryos: sequential gene expression of two neurofilament proteins in retinal ganglion cells, 65:181–189

Loh S.-H., Chan W.-T., Gong Z., Lim T.-M., Chua K.-L.
Characterization of a zebrafish (*Danio rerio*) desmin cDNA: an early molecular marker of myogenesis, 65:247–254

Lens development
Mizuno N., Mochii M., Yamamoto T.S., Takahashi T.C., Eguchi G., Okada T.S.
Pax-6 and *Prox 1* expression during lens regeneration from *Cynops* iris and *Xenopus* cornea: evidence for a genetic program common to embryonic lens development, 65:141–149

Tseng H., Matsuzaki K., Lavker R.M.
Basonuclin in murine corneal and lens epithelia correlates with cellular maturation and proliferative ability, 65:221–227

Lens regeneration
Mizuno N., Mochii M., Yamamoto T.S., Takahashi T.C., Eguchi G., Okada T.S.
Pax-6 and *Prox 1* expression during lens regeneration from *Cynops* iris and *Xenopus* cornea: evidence for a genetic program common to embryonic lens development, 65:141–149

Limbal epithelium
Tseng H., Matsuzaki K., Lavker R.M.
Basonuclin in murine corneal and lens epithelia correlates with cellular maturation and proliferative ability, 65:221–227

Low-frequency stimulation
Düsterhöft S., Putman C.T., Pette D.
Changes in FGF and FGF-receptor expression in low-frequency-stimulated rat muscles and rat satellite cell cultures, 65:203–208

Lysosomal function
Lichter-Konecki U., Moter S.E., Krawisz B.R., Schlotter M., Hipke C., Konecki D.S
Expression patterns of murine lysosome-associated membrane protein 2 (Lamp-2) transcripts during morphogenesis, 65:43–58

Lysosome-associated membrane protein-2-transcripts
Lichter-Konecki U., Moter S.E., Krawisz B.R., Schlotter M., Hipke C., Konecki D.S
Expression patterns of murine lysosome-associated membrane protein 2 (Lamp-2) transcripts during morphogenesis, 65:43–58

Mesenchyme
Benazzouz A., Duprey P.
The vimentin promoter as a tool to analyze the early events of retinoic acid-induced differentiation of cultured embryonal carcinoma cells, 65:171–180

Mesoderm
Gong S.-G., Kiba A.
The role of Xmsx-2 in the anterior-posterior patterning of the mesoderm in *Xenopus laevis*, 65:131–140

Pratt M.A.C., Crippen C.A., Ménard M.
Spontaneous retinoic acid receptor β2 expression during mesoderm differentiation of P19 murine embryonal carcinoma cells, 65:271–279

Mesothelium
Gulyás M., Dobra K., Hjerpe A.
Expression of genes coding for proteoglycans and Wilms'tumour susceptibility gene 1 (*WT1*) by variously differentiated benign human mesothelial cells, 65:89–96

Mitochondria
Joffroy S., Letellier T., Rossignol R., Malgat M., Delage J.-P., Powell J.A., Mazat J.-P., Koenig J.
Modification of mitochondrial metabolism in fibroblasts from mice with a skeletal muscle mutation (muscular dysgenesis): Evidence of embryonic communication between myoblasts and fibroblasts, 65:261–270

Monocytes
Pujol B.F., Lucibello F.C., Gehling U.M., Lindemann K., Weidner N., Zuzarte M.-L., Adamkiewicz J., Elsässer H.-P., Müller R., Havemann K.
Endothelioid cells derived from human CD14 positive monocytes, 65:287–300

Muscle development
Düsterhöft S., Pette D.
Evidence that acidic fibroblast growth factor promotes maturation of rat satellite-cell-derived myotubes in vitro, 65:161–169

Düsterhöft S., Putman C.T., Pette D.
Changes in FGF and FGF-receptor expression in low-frequency-stimulated rat muscles and rat satellite cell cultures, 65:203–208

Loh S.-H., Chan W.-T., Gong Z., Lim T.-M., Chua K.-L.
Characterization of a zebrafish (*Danio rerio*) desmin cDNA: an early molecular marker of myogenesis, 65:247–254

Low R.B., White S.L., Low E.S., Neuville P., Bochaton-Piallat M.-L., Gabbiani G.
Age dependence of smooth muscle myosin expression by cultured rat aortic smooth muscle, 65:151–159

Muscular dysgenesis
Joffroy S., Letellier T., Rossignol R., Malgat M., Delage J.-P., Powell J.A., Mazat J.-P., Koenig J.
Modification of mitochondrial metabolism in fibroblasts from mice with a skeletal muscle mutation (muscular dysgenesis): Evidence of embryonic communication between myoblasts and fibroblasts, 65:261–270

Loh S.-H., Chan W.-T., Gong Z., Lim T.-M., Chua K.-L.
Characterization of a zebrafish (*Danio rerio*) desmin cDNA: an early molecular marker of myogenesis, 65:247–254

Myosins
Low R.B., White S.L., Low E.S., Neuville P., Bochaton-Piallat M.-L., Gabbiani G.
Age dependence of smooth muscle myosin expression by cultured rat aortic smooth muscle, 65:151–159

Neural crest
Lichter-Konecki U., Moter S.E., Krawisz B.R., Schlotter M., Hipke C., Konecki D.S.
Expression patterns of murine lysosome-associated membrane protein 2 (Lamp-2) transcripts during morphogenesis, 65:43–58

Neurofilaments
Leake D., Asch W.S., Canger A.K., Schechter N.
Gefiltin in zebrafish embryos: sequential gene expression of two neurofilament proteins in retinal ganglion cells, 65:181–189

5'non-translated leader sequences
Agarwal A.K., Parrish S.N., Blumberg D.D.
Ribosomal protein gene expression is cell type specific during development in *Dictyostelium discoideum*, 65:73–88

Oxidative phosphorylation
Joffroy S., Letellier T., Rossignol R., Malgat M., Delage J.-P., Powell J.A., Mazat J.-P., Koenig J.
Modification of mitochondrial metabolism in fibroblasts from mice with a skeletal muscle mutation (muscular dysgenesis): Evidence of embryonic communication between myoblasts and fibroblasts, 65:261–270

Pancreas
Crisera C.A., Maldonado T.S., Kadison A.S., Li M., Alkasab S.L., Laongaker M.T., Gittes G.K.
Transforming growth factor-β1 in the developing pancreas: a potential regulator of exocrine differentiation, 65:255–259

Pax-6
Mizuno N., Mochii M., Yamamoto T.S., Takahashi T.C., Eguchi G., Okada T.S.
Pax-6 and *Prox 1* expression during lens regeneration from *Cynops* iris and *Xenopus* cornea: evidence for a genetic program common to embryonic lens development, 65:141–149

Polycomb-group proteins
Tomotsune D., Takihara Y., Berger J., Duhl D., Joo S., Kyba M., Shirai M., Ohta H., Matsuda Y., Honda B.M., Simon J., Shimada K., Brock H.W., Randazzo F.
A novel member of murine Polycomb-group proteins, Sex comb on midleg homolog protein, is highly conserved, and interacts with RAE28/mph1 in vitro, 65:229–239

Preadipocyte (BIP) cell line
Inoue-Murayama M., Sugimoto Y., Niimi Y., Aso H.
Type XVIII collagen is newly transcribed during bovine adipogenesis, 65:281–285

Proliferation
Flickinger R.A.
Induction of tumor cell differentiation, 65:241–245

Prostate
Aboseif S., El-Sakka A., Young P., Cunha G.
Mesenchymal reprogramming of adult human epithelial differentiation, 65:113–118

Proteases
Fehlner-Gardiner CC., Cao H., Jackson-Boeters L., Nakamura T., Elliott BE., Uniyal S., Chan BMC.
Characterization of a functional relationship between hepatocyte growth factor and mouse bone marrow-derived mast cells, 65:27–42

Proteoglycans
Gulyás M., Dobra K., Hjerpe A.
Expression of genes coding for proteoglycans and Wilms'tumour susceptibility gene 1 (*WT1*) by variously differentiated benign human mesothelial cells, 65:89–96

Lichter-Konecki U., Moter SE., Krawisz BR., Schlotter M., Hipke C., Konecki DS
Expression patterns of murine lysosome-associated membrane protein 2 (Lamp-2) transcripts during morphogenesis, 65:43–58

Prox 1
Mizuno N., Mochii M., Yamamoto TS., Takahashi TC., Eguchi G., Okada TS.
Pax-6 and *Prox 1* expression during lens regeneration from *Cynops* iris and *Xenopus* cornea: evidence for a genetic program common to embryonic lens development, 65:141–149

Quick freeze deep etch
Liang F., Kachar B., Ding M., Zhai Z., Wu X-R., Sun T-T.
Urothelial hinge as a higly specialized membrane: detergent-insolubility, urohingin association, and in vitro formation, 65:59–69

Regeneration
Leake D., Asch WS., Canger AK., Schechter N.
Gefiltin in zebrafish embryos: sequential gene expression of two neurofilament proteins in retinal ganglion cells, 65:181–189

Aboseif S., El-Sakka A., Young P., Cunha G.
Mesenchymal reprogramming of adult human epithelial differentiation, 65:113–118

Retinal ganglion cells
Leake D., Asch WS., Canger AK., Schechter N.
Gefiltin in zebrafish embryos: sequential gene expression of two neurofilament proteins in retinal ganglion cells, 65:181–189

Retinoic acid
Essner JJ., Johnson RG., Hackett PB Jr.
Overexpression of thyroid hormone receptor β1 during zebrafish embryogenesis disrupts hindbrain patterning and implicates retinoic acid receptors in the control of *hox* gene expression, 65:1–11

Benazzouz A., Duprey P.
The vimentin promoter as a tool to analyze the early events of retinoic acid-induced differentiation of cultured embryonal carcinoma cells, 65:171–180

Tomotsune D., Takihara Y., Berger J., Duhl D., Joo S., Kyba M., Shirai M., Ohta H., Matsuda Y., Honda BM., Simon J., Shimada K., Brock HW., Randazzo F.
A novel member of murine Polycomb-group proteins, Sex comb on midleg homolog protein, is highly conserved, and interacts with RAE28/mph1 in vitro, 65:229–239

Pratt M.A.C., Crippen C.A., Ménard M.
Spontaneous retinoic acid receptor β2 expression during mesoderm differentiation of P19 murine embryonal carcinoma cells, 65:271–279

Essner J.J., Johnson R.G., Hackett P.B Jr.
Overexpression of thyroid hormone receptor α1 during zebrafish embryogenesis disrupts hindbrain patterning and implicates retinoic acid receptors in the control of *hox* gene expression, 65:1–11

Ribosomal proteins
Agarwal A.K., Parrish S.N., Blumberg D.D.
Ribosomal protein gene expression is cell type specific during development in *Dictyostelium discoideum*, 65:73–88

Satellite cell culture
Düsterhöft S., Pette D.
Evidence that acidic fibroblast growth factor promotes maturation of rat satellite-cell-derived myotubes in vitro, 65:161–169
Düsterhöft S., Putman C.T., Pette D.
Changes in FGF and FGF-receptor expression in low-frequency-stimulated rat muscles and rat satellite cell cultures, 65:203–208

Seminal vesicle
Aboseif S., El-Sakka A., Young P., Cunha G.
Mesenchymal reprogramming of adult human epithelial differentiation, 65:113–118

Sex comb
Tomotsune D., Takihara Y., Berger J., Duhl D., Joo S., Kyba M., Shirai M., Ohta H., Matsuda Y., Honda B.M., Simon J., Shimada K., Brock H.W., Randazzo F.
A novel member of murine Polycomb-group proteins, Sex comb on midleg homolog protein, is highly conserved, and interacts with RAE28/mph1 in vitro, 65:229–239

Smooth muscle
Low R.B., White S.L., Low E.S., Neuville P., Bochaton-Piallat M.-L., Gabbiani G.
Age dependence of smooth muscle myosin expression by cultured rat aortic smooth muscle, 65:151–159

Stem cells
Tseng H., Matsuzaki K., Lavker R.M.
Basonuclin in murine corneal and lens epithelia correlates with cellular maturation and proliferative ability, 65:221–227

TGF-(symbol)beta
Low R.B., White S.L., Low E.S., Neuville P., Bochaton-Piallat M.-L., Gabbiani G.
Age dependence of smooth muscle myosin expression by cultured rat aortic smooth muscle, 65:151–159

Thyroid hormone receptor
Essner J.J., Johnson R.G., Hackett P.B. Jr.
Overexpression of thyroid hormone receptor α1 during zebrafish embryogenesis disrupts hindbrain patterning and implicates retinoic acid receptors in the control of *hox* gene expression, 65:1–11

Tongue
Rothnagel J.A., Seki T., Ogo M., Longley M.A., Wojcik S.M., Bundman D.S., Bickenbach J.R., Roop D.R.
The mouse keratin 6 isoforms are differentially expressed in the hair follicle, footpad, tongue and activated epidermis, 65:119–130

Transcriptional control
Tseng H., Matsuzaki K., Lavker R.M.
Basonuclin in murine corneal and lens epithelia correlates with cellular maturation and proliferative ability, 65:221–227
Flickinger R.A.
Induction of tumor cell differentiation, 65:241–245
Fahrenkrug S.C., Dahlquist M.O., Clark K.J., Hackett P.B.
Dynamic and tissue-specific expression of eIF4E during zebrafish embryogenesis, 65:191–201

Transfilter-recombination
Koike T., Yasugi S.
In Vitro analysis of mesenchymal influences on the differentiation of stomach epithelial cells of the chicken embryo. 65:13–25

Transforming-growth-factor-β1
Crisera C.A., Maldonado T.S., Kadison A.S., Li M., Alkasab S.L., Laongaker M.T., Gittes G.K.
Transforming growth factor-β1 in the developing pancreas: a potential regulator of exocrine differentiation, 65:255–259

Transgenic mice
Wojcik S.M., Imakado S., Seki T., Longley M.A., Petherbridge L., Bundman D.S., Bickenbach J.R., Rothnagel J.A., Roop D.R.
Expression of *MK6a* dominant-negative and C-terminal mutant transgenes in mice has distinct phenotypic consequences in the epidermis and hair follicle, 65:97–112
Rothnagel J.A., Seki T., Ogo M., Longley M.A., Wojcik S.M., Bundman D.S., Bickenbach J.R., Roop D.R.
The mouse keratin 6 isoforms are differentially expressed in the hair follicle, footpad, tongue and activated epidermis, 65:119–130

Translation
Fahrenkrug S.C., Dahlquist M.O., Clark K.J., Hackett P.B.
Dynamic and tissue-specific expression of eIF4E during zebrafish embryogenesis, 65:191–201
Agarwal AK., Parrish SN., Blumberg DD.
Ribosomal protein gene expression is cell type specific during development in *Dictyostelium discoideum*, 65:73–88

Uroplakin
Liang F., Kachar B., Ding M., Zhai Z., Wu X.-R., Sun T.-T.
Urothelial hinge as a higly specialized membrane: detergent-insolubility, urohingin association, and in vitro formation, 65:59–69

Urothelium
Liang F., Kachar B., Ding M., Zhai Z., Wu X.-R., Sun T.-T.
Urothelial hinge as a higly specialized membrane: detergent-insolubility, urohingin association, and in vitro formation, 65:59–69

Ventral mesoderm
Fahrenkrug SC., Dahlquist MO., Clark KJ., Hackett PB.
Dynamic and tissue-specific expression of eIF4E during zebrafish embryogenesis, 65:191–201

Vimentin
Benazzouz A., Duprey P.
The vimentin promoter as a tool to analyze the early events of retinoic acid-induced differentiation of cultured embryonal carcinoma cells, 65:171–180

Wilms' tumour susceptibility gene 1 (*WT1*)
Gulyás M., Dobra K., Hjerpe A.
Expression of genes coding for proteoglycans and Wilms'tumour susceptibility gene 1 (*WT1*) by variously differentiated benign human mesothelial cells, 65:89–96

Xenopus laevis
Gong S.-G., Kiba A.
The role of Xmsx-2 in the anterior-posterior patterning of the mesoderm in *Xenopus laevis*, 65:131–140
Mizuno N., Mochii M., Yamamoto T.S., Takahashi T.C., Eguchi G., Okada T.S.
Pax-6 and *Prox 1* expression during lens regeneration from *Cynops* iris and *Xenopus* cornea: evidence for a genetic program common to embryonic lens development, 65:141–149

Xmsx-2
Gong S.-G., Kiba A.
The role of Xmsx-2 in the anterior-posterior patterning of the mesoderm in *Xenopus laevis*, 65:131–140

Zebrafish

Essner J.J., Johnson R.G., Hackett P.B. Jr.
Overexpression of thyroid hormone receptor α1 during zebrafish embryogenesis disrupts hindbrain patterning and implicates retinoic acid receptors in the control of *hox* gene expression, 65:1–11

Leake D., Asch W.S., Canger A.K., Schechter N.
Gefiltin in zebrafish embryos: sequential gene expression of two neurofilament proteins in retinal ganglion cells, 65:181–189

Fahrenkrug S.C., Dahlquist M.O., Clark K.J., Hackett P.B.
Dynamic and tissue-specific expression of eIF4E during zebrafish embryogenesis, 65:191–201

Loh S.-H., Chan W.-T., Gong Z., Lim T.-M., Chua K.-L.
Characterization of a zebrafish (*Danio rerio*) desmin cDNA: an early molecular marker of myogenesis, 65:247–254